

Popular Article

The Silent Invader: Understanding *Anaplasma phagocytophilum* and Its Global Impact

Rahul G Kadam; Amit Raj Gupta*, Saurabh Zingare, Kruti Debanath Mandal, Vinod Kumar; and Dayanidhi Jena

Assistant professor, Unit of TVCC, Faculty of Veterinary and Animal sciences, RGSC, BHU, Varanasi

*Professor and In-Charge unit of TVCC, FVAS, RGSC, BHU

*Corresponding author: Rahul G Kadam, Email: samerahul@rediffmail.com


Introduction: A Hidden Threat in the Bite of a Tick

Imagine a quiet meadow in summer. Sheep graze, deer wander through tall grass, and hikers enjoy the sunshine. Beneath this peaceful scene, an invisible drama unfolds. A microscopic bacterium *Anaplasma phagocytophilum* travels silently inside ticks, waiting for a chance to enter the bloodstream of an unsuspecting host. Once inside, it takes control of immune cells, weakening the body's defenses and opening the door to serious illness. For decades, this pathogen was thought to trouble only livestock. Today, we know it also infects humans, horses, and dogs, making it a zoonotic threat with wide-ranging consequences.

A Brief History: From Scottish Sheep to American Patients

The story begins in 1932, when veterinarians in Scotland described a puzzling illness in sheep known as tick-borne fever (TBF). Animals developed high

fever, weakness, and sometimes died from secondary infections. Over time, similar diseases were spotted in cattle and goats across Europe. By the late 20th century, scientists realized that horses and dogs could be affected by related conditions, later renamed equine granulocytic anaplasmosis (EGA) and canine granulocytic anaplasmosis (CGA). The real shock came in 1994, when doctors in the United States diagnosed human granulocytic anaplasmosis (HGA) proof that this was no longer an animal-only disease.



Meet the Culprit: What Is *Anaplasma phagocytophilum*?

Anaplasma phagocytophilum is a tiny,

Gram-negative bacterium that can't survive outside living cells. It prefers to hide inside neutrophils a type of white blood cell that normally defends the body against invaders. Instead of being destroyed, the bacterium cleverly shuts down the cell's killing mechanisms and multiplies safely within little bubbles called morulae. This ability makes it unusual: most bacteria avoid neutrophils, but *Anaplasma* has turned them into its fortress.

The Great Masquerader: Strains and Variants

Scientists once gave different names to what they thought were separate species (*Ehrlichia phagocytophila*, *Ehrlichia equi*, etc.). Eventually, genetic studies proved they were all variants of the same organism: *A. phagocytophilum*. However, not all strains behave alike. European sheep strains differ from American human strains in severity, host preference, and geography. Some infect cattle but not humans; others infect deer but rarely cause illness in livestock. This diversity complicates diagnosis and control.

Life Cycle: The Tick Connection

Ticks of the *Ixodes* genus—the same ones that spread Lyme disease—are the primary carriers. The bacterium moves between ticks and animals in a repeating cycle. Importantly, ticks do not pass the infection to their offspring, meaning the chain relies

on feeding from infected hosts. Rodents, deer, and livestock act as reservoirs, keeping the bacterium circulating in nature. Birds may even help spread infected ticks over long distances. For humans, a single tick bite can transmit the pathogen after hours of feeding.



How the Disease Strikes

Once inside the body, *Anaplasma* causes:

- High fever and chills (often mistaken for flu)
- Drop in white blood cells (neutropenia, lymphocytopenia)
- Low platelet count (thrombocytopenia, leading to bleeding risk)
- Immune suppression, making patients vulnerable to other infections

In sheep and cattle, the consequences include abortions, pneumonia, and secondary bacterial infections. In lambs, it can lead to tick pyaemia, a crippling, often fatal complication. Humans usually recover with antibiotics, but severe cases can result in multi-organ failure, especially in the elderly or immunocompromised.

A Master of Evasion

What makes *A. phagocytophilum* so persistent? Its secret lies in antigenic variation the ability to change its surface proteins and stay one step ahead of the immune system. This means previously infected animals (or humans) may still get sick again with a new strain.

Global Impact and Economic Losses

The hidden toll of this infection is enormous. In the UK alone, tick-borne fever affects hundreds of thousands of lambs annually, leading to high mortality and reduced farm income. Dairy cattle suffer drops in milk yield. Abortions in sheep and cattle cause further economic strain. For human health, the rise of HGA cases in Europe and North America highlights the growing importance of tick-borne zoonoses in a warming world where ticks are expanding their range.

Diagnosis and Treatment

Because symptoms mimic flu, human cases are often misdiagnosed. Laboratory tests (PCR, blood smears, serology) are required for confirmation.

Treatment

Doxycycline remains the drug of choice for humans and animals. Other antibiotics show limited success. No vaccines are yet available.


Prevention: Stopping the Tick

For now, prevention focuses on:

- Tick control in livestock (dipping,

pour-on insecticides)

- Strategic use of antibiotics in at-risk herds
- Avoidance measures for people (repellents, protective clothing, careful tick checks after outdoor activity).



Looking Ahead: The Unanswered Questions

Despite decades of research, mysteries remain:

- Where does the bacterium hide between infections?
- Why do some strains infect humans while others don't?
- How exactly does it manipulate neutrophils so effectively?

Ongoing genome sequencing and molecular studies promise new insights. A safe, effective vaccine especially for livestock could one day transform control efforts.

Conclusion: The Quiet Power of a Tiny Pathogen

Anaplasma phagocytophilum is a reminder that even the smallest organisms can have vast impacts on animal health, farm economics, and human well-being. From Scottish sheep in the 1930s to American patients in the 1990s, its story illustrates

how pathogens cross boundaries between species. As climate change alters tick habitats and brings humans, livestock, and wildlife into closer contact, the risk only grows. Understanding and controlling this stealthy invader will remain a crucial challenge for both veterinary and human medicine.